

**animal**

<http://journals.cambridge.org/ANM>

Additional services for **animal**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



---

## Mid-infrared prediction of lactoferrin content in bovine milk: potential indicator of mastitis

H. Soyeurt, C. Bastin, F. G. Colinet, V. M.-R. Arnould, D. P. Berry, E. Wall, F. Dehareng, H. N. Nguyen, P. Dardenne, J. Schefers, J. Vandenplas, K. Weigel, M. Coffey, L. Théron, J. Detilleux, E. Reding, N. Gengler and S. McParland

animal / Volume 6 / Issue 11 / November 2012, pp 1830 - 1838

DOI: 10.1017/S1751731112000791, Published online:

**Link to this article:** [http://journals.cambridge.org/abstract\\_S1751731112000791](http://journals.cambridge.org/abstract_S1751731112000791)

### How to cite this article:

H. Soyeurt, C. Bastin, F. G. Colinet, V. M.-R. Arnould, D. P. Berry, E. Wall, F. Dehareng, H. N. Nguyen, P. Dardenne, J. Schefers, J. Vandenplas, K. Weigel, M. Coffey, L. Théron, J. Detilleux, E. Reding, N. Gengler and S. McParland (2012). Mid-infrared prediction of lactoferrin content in bovine milk: potential indicator of mastitis. *animal*, 6, pp 1830-1838 doi:10.1017/S1751731112000791

**Request Permissions :** [Click here](#)

## Mid-infrared prediction of lactoferrin content in bovine milk: potential indicator of mastitis

H. Soyeurt<sup>1,2†</sup>, C. Bastin<sup>1</sup>, F. G. Colinet<sup>1</sup>, V. M.-R. Arnould<sup>1,3</sup>, D. P. Berry<sup>4</sup>, E. Wall<sup>5</sup>, F. Dehareng<sup>6</sup>, H. N. Nguyen<sup>6</sup>, P. Dardenne<sup>6</sup>, J. Scheifers<sup>7</sup>, J. Vandenplas<sup>1,2</sup>, K. Weigel<sup>7</sup>, M. Coffey<sup>5</sup>, L. Théron<sup>8</sup>, J. Detilleux<sup>8</sup>, E. Reding<sup>9</sup>, N. Gengler<sup>1,2</sup> and S. McParland<sup>4</sup>

<sup>1</sup>Animal Science Unit, Gembloux Agro-Bio Tech, University of Liège, 5030 Gembloux, Belgium; <sup>2</sup>National Fund for Scientific Research, B-1000 Brussels, Belgium; <sup>3</sup>Convis Herdbuch, Zone artisanale et commerciale no 4, 9085 Ettelbruck, Luxembourg; <sup>4</sup>Animal and Grassland Research and Innovation Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland; <sup>5</sup>Sustainable Livestock Systems Group, Scottish Agricultural College, Easter Bush Campus, Midlothian EH25 9RG, UK; <sup>6</sup>Valorisation of Agricultural Products Department, Walloon Research Centre, 5030 Gembloux, Belgium; <sup>7</sup>Department of Dairy Science, University of Wisconsin, 1675 Observatory Drive, Madison, WI 53706, USA; <sup>8</sup>Clinical Department for Production Animals – Ruminants Clinic, Faculty of Veterinary Medicine, University of Liège, 4000 Liège, Belgium; <sup>9</sup>Walloon Breeding Association, 5590 Ciney, Belgium

(Received 16 September 2011; Accepted 27 January 2012; First published online 27 April 2012)

Lactoferrin (LTF) is a milk glycoprotein favorably associated with the immune system of dairy cows. Somatic cell count is often used as an indicator of mastitis in dairy cows, but knowledge on the milk LTF content could aid in mastitis detection. An inexpensive, rapid and robust method to predict milk LTF is required. The aim of this study was to develop an equation to quantify the LTF content in bovine milk using mid-infrared (MIR) spectrometry. LTF was quantified by enzyme-linked immunosorbent assay (ELISA), and all milk samples were analyzed by MIR. After discarding samples with a coefficient of variation between 2 ELISA measurements of more than 5% and the spectral outliers, the calibration set consisted of 2499 samples from Belgium ( $n = 110$ ), Ireland ( $n = 1658$ ) and Scotland ( $n = 731$ ). Six statistical methods were evaluated to develop the LTF equation. The best method yielded a cross-validation coefficient of determination for LTF of 0.71 and a cross-validation standard error of 50.55 mg/l of milk. An external validation was undertaken using an additional dataset containing 274 Walloon samples. The validation coefficient of determination was 0.60. To assess the usefulness of the MIR predicted LTF, four logistic regressions using somatic cell score (SCS) and MIR LTF were developed to predict the presence of mastitis. The dataset used to build the logistic regressions consisted of 275 mastitis records and 13 507 MIR data collected in 18 Walloon herds. The LTF and the interaction  $SCS \times LTF$  effects were significant ( $P < 0.001$  and  $P = 0.02$ , respectively). When only the predicted LTF was included in the model, the prediction of the presence of mastitis was not accurate despite a moderate correlation between SCS and LTF ( $r = 0.54$ ). The specificity and the sensitivity of models were assessed using Walloon data (i.e. internal validation) and data collected from a research herd at the University of Wisconsin – Madison (i.e. 5886 Wisconsin MIR records related to 93 mastitis events – external validation). Model specificity was better when LTF was included in the regression along with SCS when compared with SCS alone. Correct classification of non-mastitis records was 95.44% and 92.05% from Wisconsin and Walloon data, respectively. The same conclusion was formulated from the Hosmer and Lemeshow test. In conclusion, this study confirms the possibility to quantify an LTF indicator from milk MIR spectra. It suggests the usefulness of this indicator associated to SCS to detect the presence of mastitis. Moreover, the knowledge of milk LTF could also improve the milk nutritional quality.

**Keywords:** cow, lactoferrin, milk, mid-infrared spectrometry, mastitis

### Implications

Lactoferrin (LTF) is a glycoprotein naturally present in milk with known favorable associations with the immune system. The quantification of this molecule directly in milk could be useful in the development of animal breeding programs to

improve mastitis resistance of dairy cows and to improve the nutritional quality of bovine milk. To achieve these objectives an inexpensive, fast and robust method to quantify LTF directly from bovine milk is required. The results of this study show the potential of mid-infrared (MIR) spectrometry to predict LTF. MIR spectrometry is the method of choice worldwide to routinely predict milk components. Thus, the equation developed could potentially be exploited at low

† E-mail: hsoyeurt@ulg.ac.be

cost in the short term. Moreover, this study showed also that the use of predicted LTF and somatic cell score (SCS) slightly improved the possibility to detect the presence of clinical mastitis in dairy cows over and above using SCS alone.

## Introduction

Lactoferrin (LTF) is an iron-binding glycoprotein naturally present in milk and secreted mainly by the mammary epithelial cells. Its concentration is higher during the later lactation stages (0.25 to 0.40 g/l) compared with early lactation stages (Hagiwara *et al.*, 2003; Baumrucker *et al.*, 2005; Pugovel *et al.*, 2005). LTF can also be released by polymorphonuclear neutrophils during inflammation (Kuttila *et al.*, 2004).

LTF belongs to the transferrin gene family (Mead and Tweedie, 1990; Ward *et al.*, 2005) and has reverse iron-binding properties mainly explaining the different roles allotted to LTF in the regulation of the immune system (Mead and Tweedie, 1990; Baker and Baker, 2004). LTF seems to act as a general anti-bacterial and antifungal molecule (Baker, 2005; Farnaud and Evans, 2003), to be involved in host defence mechanisms (Baveye *et al.*, 1999; Baumrucker, 2000; Ward *et al.*, 2005), and to modulate the inflammatory process (Farnaud and Evans, 2003). Therefore, the content of LTF in bovine milk may have potential as an indicator of the presence of mastitis and could be used in an animal breeding program to augment the accuracy of selection for mastitis resistance in dairy cows. Indeed, high LTF concentration in milk (2.3 g/l of milk) compared with the expected normal level (0.1 to 0.4 g/l of milk) produced by a specific cow may indicate clinical or subclinical mastitis (Kuttila *et al.*, 2004). Moreover, naturally increasing the content of LTF in milk through genetic selection could also have beneficial effects on human health (Tsuda *et al.*, 2000).

For practical reasons (i.e. analysis time, skilled staff, analysis cost), the current methods used to quantify the LTF content in milk such as enzyme-linked immunosorbent assay (ELISA) or the immunodiffusion method are not feasible for a large number of samples as is necessary for accurate estimation of breeding values. Therefore, an alternative method that is rapid and low cost is required.

Recent research has shown the potential of mid-infrared (MIR) spectrometry to predict components in milk, for example, milk fat composition, in addition to the routine predictions of milk fat, protein, lactose and urea undertaken globally (Rutten *et al.*, 2006; Soyeurt *et al.* 2006 and 2011). Similarly, MIR indicators of protein fractions such as casein and lactoglobulin (Barbano and Dellavalle, 1987; Sorensen *et al.*, 2003) have been obtained. As it is possible to determine the total content of protein and some specific milk proteins, it could be possible to predict the LTF content of milk using MIR as LTF is a natural milk glycoprotein. This hypothesis was tested by Soyeurt *et al.* (2007) who suggested the possibility of quantification of an indicator of LTF in bovine milk using MIR spectrometry with a cross-validation coefficient of determination ( $R_{CV}^2$ ) of 0.75. However, the study of Soyeurt *et al.* (2007) had several limitations; only 69 samples collected in the Walloon region of Belgium were

used to develop the equation through a partial least squares regression (PLS). Recent studies to predict other components in milk have shown a benefit in the accuracy of calibration equations with (1) more complex calibration procedures, (2) a larger number of samples from a wider range of production systems and (3) greater animal diversity (Dal Zotto *et al.*, 2008; Soyeurt *et al.*, 2011).

The first aim of the current study was to develop a calibration equation to quantify the LTF content in milk using a large number of samples collected in Belgium, Ireland and Scotland from cows of different breeds and from different production systems. Pre-treatments applied to the spectral data were tested to improve the robustness of the MIR prediction of LTF content. The second aim was to quantify the usefulness of this LTF prediction to identify the presence of clinical mastitis.

## Material and methods

### *Spectral and LTF data used in the calibration set*

Milk samples (40 ml each in Ireland and Scotland and 2 × 60 ml each in Belgium (i.e. one for the MIR analysis and one for the LTF quantification)) were collected between April 2005 and April 2006 in Belgium, between April 2009 and August 2009 in Ireland and in August 2009 in Scotland. All samples were analyzed using MilkoScan FT6000 spectrometers (Foss, Hillerod, Denmark) that outputs one spectrum of 1060 data points in the IR range from 900/cm to 5000/cm for each milk sample analyzed. Belgian milk samples were analyzed by MIR in the milk laboratory 'Comité du lait' (Battice, Belgium) and the remaining samples were analyzed at Teagasc Moorepark (Fermoy, Co. Cork, Ireland).

Samples were preserved with bronopol and the LTF concentration in whole milk samples was measured at least in duplicate with a commercial ELISA kit (Bovine Lactoferrin ELISA Quantification Kit, Bethyl Laboratories Inc., Montgomery, TX, USA). The procedure was carried out according to the manufacturer's instructions. The samples were diluted 1:1000; 1:2000; 1:4000; 1:6000; 1:8000 or 1:10 000 in sample buffer. The LTF concentrations used for the calibration were the average of at least two measures taken on the same milk sample. Only samples with a coefficient of variation between two repeated measurements of LTF quantification of <5% were used to develop the calibration equation. For Irish and Scottish samples, ELISA analysis was undertaken by Enfer Laboratories (Naas Co. Kildare, Ireland). Belgian milk samples were analyzed by Gembloux Agro-Bio Tech – University of Liège (Gembloux, Belgium).

A principal component analysis was undertaken across all spectral data included in the calibration set and the standardized Mahalanobis distance was calculated. As described by Williams (2007), a spectral outlier is a spectrum which differs from the mean of the population by 3 or more standardized Mahalanobis distance. Using this threshold, 60 samples were deemed to be outliers and were discarded. Therefore, the final calibration dataset included 110 Walloon samples and 1658 Irish and 731 Scottish samples.

The Walloon samples (i.e. 110 records) were selected to maximize the spectral variability from the 1609 samples collected in a previous study (Soyeurt *et al.*, 2007) from six different breeds (dual-purpose Belgian Blue, Holstein–Friesian, Jersey, Montbeliarde, Normande, non-Holstein Meuse-Rhine-Yssel type Red and White) across seven herds. All of these samples were obtained during the routine milk recording managed by the Walloon Breeding Association (AWE, Ciney, Belgium) and frozen at  $-26^{\circ}\text{C}$ . More details are given by Soyeurt *et al.* (2007).

Irish and Scottish samples were analyzed fresh one day after collection. The cows sampled in the Irish research herds were fed predominantly grazed grass and included animals of different breeds (Holstein–Friesian, Holstein, Friesian, Jersey, crossbred Jersey and Holstein–Friesian, Montbeliarde, Normande, Norwegian Red, crossbred Norwegian Red and Holstein Friesian; Prendiville *et al.*, 2010) and different strains of Holstein–Friesian (Coleman *et al.*, 2009). Cows from Scotland were from two genetically divergent lines (divergent for milk solids) and were fed two different diets (Coffey *et al.*, 2004). Approximately half were fed a predominantly forage only diet and the other half a diet consisting of  $\sim 60\%$  silage and 40% concentrate.

#### Calibration models

A PLS approach was undertaken to relate the ELISA LTF values to the spectral data using WINISI III software (<http://www.winisi.com/>; Foss, Hillerod, Denmark). This regression technique requires cross-validation to prevent over-fitting to the data. Cross-validation obtains validation errors by partitioning the calibration set into several groups (50 groups were used in this study). The calibration is performed on the dataset with one group excluded and the equations developed are applied to the excluded group; this is iterated with a different group excluded at each iteration. The residuals from the prediction for each sample were combined into a standard error of cross-validation (Sinnaeve *et al.*, 1994). In order to assess the efficiency of the calibration equation, different statistical parameters were estimated: mean LTF ELISA content, standard deviation of ELISA LTF content (s.d.), standard error of calibration (s.e.c.), calibration coefficient of determination ( $R_c^2$ ), standard error of cross-validation (SECV) and cross-validation coefficient of determination ( $R_{cv}^2$ ). The ratio of SECV to s.d. (ratio of s.e. prediction to s.d.; RPD) was also calculated (William and Norris, 2001) in order to estimate the efficiency of calibration.

An equation developed for one spectrometer could provide results slightly biased on another instrument. To improve the accuracy of the MIR prediction across instruments, a 'repeatability file' was generated by recording the MIR spectrum of several milk samples provided by five different MilkoScan FT6000 spectrometers (Foss) and used in the calibration process based on a methodology proposed by Westerhaus (1990). More details about the repeatability file were provided by Soyeurt *et al.* (2011).

Generating the derivative of the spectral data permits a 'sharpening' of the absorption bands. First and second derivatives were obtained using the following formula:

$$dx_k = x_{k - \frac{g-1}{2}} - x_{k + \frac{g-1}{2}}$$

where  $dx_k$  is the value of the derivative for the spectral data point  $k$ ;  $x$  is the absorbance;  $g$  is an odd integer strictly positive entitled 'gap' (i.e. five consecutive spectral data points in this study; Hruschka, 1987). The second derivative used the absorbance obtained for the first derivative instead of the absorbance of spectra. Six different statistical approaches were evaluated to develop the prediction equations:

- method 1: PLS and no pre-treatment on the spectral data;
- method 2: PLS + the use of a repeatability file;
- method 3: PLS + the use of a first derivative pre-treatment on the spectral data;
- method 4: PLS + the use of a first derivative pre-treatment + repeatability file;
- method 5: PLS + the use of a second derivative pre-treatment;
- method 6: PLS + the use of a second derivative pre-treatment + repeatability file.

A T-outlier test was used to remove potential outliers from the reference ELISA data (Winisi software, Foss, Hillerod, Denmark). The critical T-outlier value was set to 2.5 (i.e. recommended default value). If the difference between the predicted and reference LTF values was  $>2.5$  times the SEC, the sample was considered to be an outlier and deleted. This methodology was iterated twice.

#### External validation

An external validation was conducted using 274 samples collected from 14 Walloon herds in June and July 2009, and again during January and May 2011. Samples were collected from different breeds: dual-purpose Belgian Blue, Holstein–Friesian, Montbeliarde and non-Holstein Meuse-Rhine-Yssel type Red and White. The developed equations were applied to this dataset and the validation coefficient of determination was calculated ( $R_v^2$ ).

#### Prediction of clinical mastitis

To assess the usefulness of the LTF MIR prediction to detect mastitis, the LTF calibration equation was applied to the recorded spectral data from cows with associated mastitis information.

**Calibration set.** Mastitis records were collected from 18 farms located in the Walloon region of Belgium between January 2007 and the beginning of August 2011 (two farms from 2007, two farms from 2008, two farms from 2009, five farms from 2010 and seven farms from 2011). A total of 1750 mastitis events were recorded from 864 dairy cows. The average incidence of mastitis for these herds was 30.17% with an s.d. of 15.30%. The minimum mastitis incidence was 9.09% and the maximum was 78.38%. These farms were involved in the Walloon milk recording program; therefore, the composition of milk from all cows in lactation in the herds were analyzed every 4 to 6 weeks using a Foss MilkoScan FT6000 (Foss). Spectral data and somatic cell count (SCC) were recorded. LTF content was estimated by applying the LTF MIR equation developed in this study on the recorded spectra. The somatic cell score (SCS) was obtained by the following formula:  $\text{SCS} = (\log_2(\text{SCC}/100\,000)) + 3$ . SCS was calculated because its distribution is more normal compared with the distribution of SCC data. The mastitis events were merged with the MIR records based on the

date of the mastitis event  $\pm 7$  days. Only Holstein cows were considered in this study. Finally, 275 mastitis events had an associated SCS and MIR spectral records. Therefore, the final dataset contained 13 507 MIR records with 275 mastitis events from 1564 Holstein cows.

Logistic regression (PROC LOGISTIC in SAS; SAS Institute, 1999) was used to develop an equation to detect the presence of mastitis (1 = mastitis; 0 = no mastitis). Independent variables included in the model were SCS, predicted MIR LTF and their interaction. To avoid bias from correlated records among cows, the generalized estimating equations (PROC GENMOD in SAS; SAS Institute, 1999) were tested but the results were similar and sometimes worse than those observed for PROC LOGISTIC. This could be explained by the large number of cows used in this study. Therefore, only the PROC LOGISTIC results will be presented in this study.

There were a small number of mastitis cases relative to non-mastitis samples; thus, a second, more 'balanced' dataset was created: only test-day MIR records from all cows within  $\pm 60$  days of the 275 mastitis cases were kept. This dataset contained 677 records.

Logistic regression models were built from this dataset. Comparisons were made based on different statistical parameters such as the deviance and Pearson's Goodness-of-Fit, the percent of concordant, discordant and tied samples, and the receiver operator curve (ROC) area. The significance of the different effects included in the model was assessed using the Wald test.

**Internal validation.** The internal validation of the equations (i.e. nearly all samples used for the validation were used for the calibration) was based on the  $P$  of the Hosmer and Lemeshow Goodness-of-Fit test executed by using PROC LOGISTIC in SAS with the LACKFIT option (SAS Institute, 1999). On the other hand, the coefficients of the logistic equation developed previously were applied to the entire available Walloon dataset (i.e. 13 507 MIR records related to 275 mastitis events) in order to predict the logit value, which is an estimation of the  $P$  of the presence of mastitis. The thresholds used were the following:

Predicted logit value = 0, then predicted mastitis event was coded as 0.

Predicted logit value  $> 0$ , then predicted mastitis event was coded as 1.

Predicted logit value  $< 0$ , then predicted mastitis event was coded as 0.

Then, the specificity (i.e. the ability of the model to predict the absence of mastitis) and the sensitivity (i.e. the ability of the model to predict the presence of mastitis) were calculated by the following formulas:

Sensitivity =

$$\frac{\sum \text{Samples correctly classified as provided by cows with mastitis}}{\sum \text{Samples provided by cows with mastitis}}$$

Sensitivity =

$$\frac{\sum \text{Samples correctly classified as provided by healthy cows}}{\sum \text{Samples provided by healthy cows}}$$

**External validation.** A total of 5886 milk samples from 800 Holstein cows were collected in the research farm of the University of Wisconsin – Madison. All samples were analyzed by MIR spectrometry by the milk lab 'AgSource Cooperative Service' (Menomonie, WI, USA) using a MilkoScan FT6000 (Foss) between January 2009 and February 2011. The spectral data and SCC were recorded. SCS was calculated by using the formula shown previously. LTF content was estimated using the developed MIR calibration equation. Moreover, 540 mastitis events were recorded between January 2009 and December 2010. The mastitis incidence was 25.39% for 2009 and 32.32% for 2010. By using the same approach of data edits used in the Walloon dataset (i.e. date of mastitis event  $\pm 7$  days), only 93 events had an associated MIR spectral record. The same methodology as the one explained for the internal validation using all Walloon dataset was used in order to calculate the  $P$  for the presence of mastitis in the external validation dataset and therefore, to evaluate the specificity and the sensitivity of the model.

## Results and discussion

### LTF calibration equations


The calibration dataset used contained 2499 samples and had a mean and s.d. for the LTF content of  $163.29 \pm 103.40$  mg/l of milk. The robustness of the six approaches to derive the equations is shown in Table 1. Because of the use of the T-outlier test, the number of samples actually used to develop the different equations differed. However, the maximum number of samples discarded by this test was 61 (i.e.  $2499 - 2438 = 61$ ), which represents  $< 2.50\%$  of the total amount of samples. Across the six different statistical approaches evaluated, the  $R_{CV}^2$  ranged from 0.69 to 0.73. Clearly, the equation combining a first derivative pre-treatment and the use of the repeatability file gave the most accurate results (i.e. high  $R_{CV}^2$  (0.72) and RPD (1.86); highest  $R_v^2$  (0.60)). The SECV was 50.55 mg/l of milk. Soyeyrt *et al.* (2011) also concluded that the use of the first-derivative and the repeatability files resulted in the most accurate prediction of milk fatty acids.

The increase in  $R_{CV}^2$  and RPD in the method including both the first derivative and the repeatability file compared with the method including only a first derivative can be explained by the

**Table 1** Statistical parameters for the six methods used to develop the lactoferrin equation

	$n$	$R_c^2$	$R_{CV}^2$	RPD	$R_v^2$
No pre-treatment	2445	0.71	0.70	1.83	0.29
First derivative	2463	0.74	0.73	1.91	0.43
First derivative + repeatability file	2442	0.72	0.71	1.86	0.60
Second derivative	2459	0.73	0.72	1.90	0.53
Second derivative + repeatability file	2438	0.70	0.69	1.81	0.51
Repeatability file	2445	0.69	0.69	1.79	0.27

$R_c^2$  = calibration coefficient of determination;  $R_{CV}^2$  = cross-validation coefficient of determination; RPD = ratio of s.d. of the reference enzyme-linked immunosorbent assay values to the standard error of cross-validation;  $R_v^2$  = validation coefficient of determination estimated from 274 Walloon samples.



**Figure 1** Distribution of lactoferrin enzyme-linked immunosorbent assay (ELISA) values (mg/l of milk).

difference in the number of samples originating from Ireland and Scotland and those originating from Belgium. On one hand, the validation dataset contained only Walloon milk samples. In contrast, the calibration dataset contained a very low number of Walloon samples relative to Scottish and Irish samples. Therefore, because of the large number of Walloon samples in the repeatability file, the use of this file during the calibration process allowed us to better consider the spectral variability present in the Walloon region of Belgium (more details on the repeatability file in Soyeurt *et al.*, 2011). In other words, the spectral variability of Walloon samples was not just based on 110 samples included in the calibration set but also from samples included in the repeatability file. In conclusion, the equation, which used the repeatability file, was more adapted for Walloon data as depicted by the validation results.

RPD for the prediction equation including both the first derivative and repeatability file was close to 2. If the RPD ratio is larger than 2, the calibration equation is considered good and can be used for practical purposes (Sinnavee *et al.*, 1994).

Without using any pre-treatment, the  $R_{CV}^2$  was equal to 0.70 (Table 1). Soyeurt *et al.* (2007) observed a  $R_{CV}^2$  of LTF of 0.75 using 69 milk samples. The slight difference compared with the present study can be explained by the difference in the number of samples included in the calibration set (2499 *v.* 69 samples) and the average content of LTF and the s.d. ( $163.29 \pm 103.40$  *v.*  $253.72 \pm 206.37$  mg/l of milk). This variability of LTF can be explained by the introduction of samples originating from different countries. This means that the current calibration set has a larger proportion of samples with a low content of LTF, which have a different spectral pattern (Figure 1). Soyeurt *et al.* (2006) and Rutten *et al.* (2009) showed both that  $R_c^2$  and  $R_{CV}^2$  are influenced by the content of the studied traits in milk. Therefore, the larger amount of milk samples with low LTF content in this study could negatively influence the obtained  $R_{CV}^2$ . However, the SECV obtained in this study was lower than the SECV reported by Soyeurt *et al.* (2007;

50.55 *v.* 103.93 mg/l of milk), suggesting a better accuracy of the LTF MIR prediction in this study.

Even if the  $R_{CV}^2$  and  $R_V^2$  values were different (Table 1), SECV and standard error of prediction (SEP) obtained from the method including both first-derivative and repeatability files were relatively similar (50.55 and 58.98 mg/l of milk), suggesting the development of a robust equation. In the same way, SEC and SECV were also similar. SEC was 49.90 mg/l of milk showing also the robustness of the developed equation.

The calibration dataset contained samples from three different countries. Irish and Scottish samples were analyzed by the same spectrometer located at Teagasc Moorepark (Fermoy, Co. Cork, Ireland) and the same laboratory for LTF. Belgian samples were analyzed by the milk lab 'Comité du Lait' for MIR spectrometry and by Gembloux Agro-Bio Tech for the ELISA analysis. This potential source of variation for the spectral and reference ELISA data generates an interest to combine all of the available samples into one calibration dataset for testing. When the calibration dataset contained only the Walloon samples (i.e. 110 samples), the  $R_{CV}^2$  was 0.62 with a SECV of 52.34 mg/l of milk. In the same way with the Irish and Scottish samples (i.e. 2389 samples; these samples were combined because they came from the same spectrometer and analyzed by ELISA in the same lab), the equation showed a  $R_{CV}^2$  of 0.70 with SECV of 50.93 mg/l of milk. For these equations when the first derivative was analyzed and the repeatability file included, the  $R_{CV}^2$  was lower and the SECV was greater. Therefore, the inclusion of samples coming from different country/region in one calibration dataset as undertaken in this study improved the robustness of the developed LTF calibration equation. This conclusion was substantiated by the results from the external validation. Indeed, when the Walloon validation samples were predicted using the equation developed from only Walloon samples (i.e. 110 samples), the  $R_V^2$  was 0.40 with a SEP of 77.26 mg/l of milk. When the LTF prediction of these


same validation samples was computed using the equation including Irish and Scottish samples, the  $R_v^2$  was 0.57 with a SEP of 59.15 mg/l of milk. The  $R_v^2$  obtained from the equation combining all available samples with a repeatability of ELISA values  $\leq 5\%$  was 0.60 (Table 1). Therefore, considering all data in the same calibration set improved the robustness of the LTF calibration equation.

Figure 2 shows the coefficients of the LTF equation including the repeatability file and the first derivative pre-treatment. In order to compare the coefficients between them, each coefficient was multiplied by the mean value of the corresponding spectral data. This permits accounting for the differences in the absorption of IR rays between spectral data. The three MIR regions considered were relevant. The region surrounding 1200/cm was associated with LTF and this region is known to be related to C–O bonds (Sivakesava and Irudayaraj, 2002). The region located around 1300/cm and related to COOH was also useful in the prediction of LTF. A large negative peak was observed around 1450/cm and another interesting region was located between 1700/cm and 1800/cm. Sivakesava and Irudayaraj (2002) mentioned

that the region located between 1700 and 1500/cm is related to protein bands. The range 1450 to 1200 is assumed to denote carboxylic groups of protein. Finally, the MIR data between 2500/cm and 2800/cm were also associated with LTF and this region is known to be related to lipids because it is the region for the ester linkage and C–H stretch group (Sivakesava and Irudayaraj, 2002).

#### *Mastitis indicator*

Descriptive statistics for the Walloon and Wisconsin data used to study the relationship between mastitis and predicted LTF are described in Table 2. On the basis of the skewness and kurtosis values, SCS as well as the predicted LTF approached a normal distribution. The correlation between these traits was 0.54 from the Wisconsin (i.e. 5886 MIR samples) and Walloon (i.e. 13 507 MIR records) data. As SCS is influenced by the presence of mastitis (Emanuelson *et al.*, 1988; Harmon, 1994; Pösö and Mäntysaari, 1996; Hagiwara *et al.*, 2003), we can hypothesize that predicted LTF may also be related to the presence of mastitis. The relationship between clinical or subclinical mastitis and the



**Figure 2** Coefficients of the lactoferrin mid-infrared equation multiplied by the mean of the corresponding spectral data.

**Table 2** Descriptive statistics of the dataset used to develop and validate the logistic regressions

	Mean	s.d.	Skewness	Kurtosis
Wisconsin data ( $n = 5886$ )				
SCC ( $\times 1000$ )	211.71	560.30	7.63	77.37
SCS	2.59	1.84	0.67	0.33
Predicted lactoferrin (mg/l of milk)	185.07	105.04	1.31	4.39
Walloon data ( $n = 13\ 507$ )				
SCC ( $\times 1000$ )	339.46	895.13	8.08	94.54
SCS	3.17	1.95	0.57	0.10
Predicted lactoferrin (mg/l of milk)	207.57	122.81	1.44	5.45

SCC = somatic cell count; SCS = somatic cell score.

**Table 3** Statistical parameters of the developed logistic regressions including SCS or SCS and lactoferrin

	Dataset (269/408)		
	SCS	SCS + lactof.	SCS + lactof. + SCS × lactof.
Deviance	$P < 0.01$	$P < 0.01$	$P < 0.01$
Pearson	$P = 0.60$	$P = 0.41$	$P = 0.48$
AIC	817.62	818.27	813.47
$R^2$	0.18	0.18	0.19
Wald	$P < 0.0001$	$P < 0.0001$	$P < 0.0001$
Intercept	$P < 0.0001$	$P < 0.0001$	$P = 0.0008$
SCS	$P < 0.0001$	$P < 0.0001$	$P = 0.0011$
Predicted lactoferrin		$P = 0.2425$	$P = 0.0060$
SCS × predicted lactoferrin			$P = 0.0175$
ROC area (%)	71.2	71.4	71.5
% concordant	71.0	71.3	71.4
% discordant	28.7	28.4	28.3
% tied	0.30	0.30	0.30
Hosmer and Lemeshow*	$P = 0.38$	$P = 0.17$	$P = 0.58$

SCS = somatic cells score; lactof. = prediction lactoferrin content using the developed MIR equation; SCS × lactof. = interaction between SCS and lactof.; AIC = Akaike's information criterion;  $R^2$  = coefficient of determination; ROC = receiver operator curve.

\* $P$ -value close to 1 is expected.

milk LTF content was mentioned previously by Gaunt *et al.* (1980), Hagiwara *et al.* (2003) and Kutila *et al.* (2004).

The logistic regression including only MIR predicted LTF did not reveal the presence of mastitis (the  $P$ -value of the Wald test was of 0.17).

Table 3 presents the statistical parameters of the logistic regression models including either SCS alone; both SCS and MIR predicted LTF; or SCS, LTF and their interaction (SCS × LTF). The  $P$ -values for the deviance Goodness-of-Fit test were similar between the three studied regressions. However, the  $P$ -value for the Pearson Goodness-of-Fit was lower for the logistic regression including both SCS and LTF. Akaike's information criterion was also lower for the regression including SCS, LTF and their interaction. The inclusion of LTF with the interaction SCS × LTF in combination with SCS seemed to improve slightly the explanation of the observed variability for the mastitis information because the  $R^2$  was higher for this regression (0.19). The inclusion of LTF and the interaction SCS × LTF in the model seems to be relevant because all of these fixed effects were significant based on the Wald test. The ability of the model to predict the presence of mastitis was better for the model including SCS, LTF and the interaction SCS × LTF. The ROC area and the percent of concordant samples were slightly higher for this third regression. In the same way, the percent of discordant samples was slightly lower.

The  $P$ -value for the Hosmer and Lemeshow Goodness-of-Fit were higher for the model including SCS, LTF and their interaction, suggesting a better ability of this model to predict the presence of mastitis. This observation can be explained partly by a relationship between the LTF content and SCC and their different reactions to the pathogens affecting the udder. Lindmark-Mansson *et al.* (2006) showed the link existing between LTF and SCC but also mentioned that the LTF content in milk begins to increase when the

SCC is >5000 cells/ml. Hagiwara *et al.* (2003) demonstrated that the changes in the LTF content in milk are different following the pathogens involving the mastitis in the udder.

Table 4 shows the percentage of samples correctly and incorrectly classified as well as the specificity and the sensitivity of the developed logistic regressions based on the entire Walloon dataset (i.e. 13 507 records – internal validation because some samples were used to create the regressions) and the Wisconsin dataset (i.e. 13 507 records – external validation because samples were not used to build the logistic regressions). The validation results gave different conclusions compared with the results obtained from the calibration process (Table 3). No large differences appeared between regressions. However, a slightly better sensitivity was obtained for the equation including only SCS from the Walloon data, and the one including SCS and LTF from the Wisconsin data. However, from the Walloon dataset, the difference of the sensitivity value between the equation including only SCS and the one containing SCS and LTF was slight (49.45 v. 49.09). A better specificity was observed for the model including LTF, SCS and the interaction SCS × LTF. The difference between the calibration and the validation processes can be explained mainly by: (1) the difference in the proportion of mastitis and no-mastitis events in the calibration and validation datasets and (2) the higher uncertainty for the 0 score (i.e. no-mastitis event) because maybe the model was predicting subclinical mastitis. Therefore, on the basis of the Hosmer and Lemeshow test, the existing interaction between LTF and SCC mentioned in the literature, and the specificity results, the best models in this study should be the one including SCS, LTF and their interaction.

Even though a positive correlation was observed between SCS and LTF, predicted LTF alone was not associated with the presence of mastitis. However, the knowledge of both SCS and LTF seems to slightly improve the mastitis prediction.



**Table 4** Internal and external validation of the developed logistic regressions

	Walloon data (internal validation)		Wisconsin data (external validation)	
	Mastitic cow (1)	Healthy cow (0)	Mastitic cow (1)	Healthy cow (0)
<b>Model with SCS</b>				
Predicted as 1	136	1253	32	298
Predicted as 0	139	11 979	61	5495
Specificity		90.53		94.86
Sensitivity		49.45		34.41
<b>Model with SCS and lactoferrin</b>				
Predicted as 1	135	1206	35	309
Predicted as 0	140	12 026	58	5484
Specificity		90.88		94.67
Sensitivity		49.09		37.63
<b>Model with SCS, lactoferrin and their interaction</b>				
Predicted as 1	128	1052	30	264
Predicted as 0	147	12 180	63	5529
Specificity		92.05		95.44
Sensitivity		46.54		32.26

SCS = somatic cells score; Mastitis 1 = cow having observed mastitis; Mastitis 0 = healthy cow; Prediction = 1 means mastitis and 0 means healthy cow.

Moreover, the knowledge of LTF content in milk is also important to characterize the nutritional quality of milk (Tsuda *et al.*, 2000). As many milk recording organizations begin to think about the possibility of retaining the spectral information, the prediction of LTF will be as easily available as the SCS data and could improve the detection of mastitis or the efficacy of breeding programs that aim to improve the mastitis resistance of dairy cows. On the basis of Hagiwara *et al.* (2003), a cow producing milk with <200 mg/l of LTF in milk could not inhibit the growth of *Escherichia coli*. A total of 52% of the cows studied by these authors produced milk with LTF content <200 mg/l. Therefore, the aim of a potential breeding program to improve mastitis resistance could ensure this limit of 200 µg of LTF/ml of milk. This would also be beneficial for human health and is possible because MIR LTF trait is heritable (Soyeurt *et al.*, 2007). However, before discussing the breeding objective, a lot of work is needed notably to study the relationship between MIR LTF and other common milk production traits (i.e. milk yield, fat, protein, etc.) and economic traits (i.e. fertility, longevity, etc.).

## Conclusions

This study confirms the possible computation of an indicator of LTF content in milk by MIR spectrometry. On the basis of  $R_{CV}^2$ , RPD and  $R_v^2$ , the calibration equation combining a first derivative pre-treatment as well as the use of a repeatability file is the best option to measure the LTF content in bovine milk. The slight difference between SECV and SEP suggests that the developed LTF equation is robust. Even if this prediction used alone does not improve the detection of mastitis, its combination with SCS is promising. A larger external validation exercise based on a dataset containing new mastitis information is required to confirm the specificity and sensitivity of the developed model. In conclusion, based on the validation results,

if a large additional cost for the measurement of LTF by MIR is imposed on the customers, the improvement of mastitis detection by adding LTF information may not be justified. However, as many milk recording organizations begin to retain the spectral information, the LTF prediction will be easily obtained at low cost, and therefore the combination of SCS and LTF to improve the detection of mastitis or the mastitis resistance by specific breeding programs will be interesting. The introduction of LTF in a breeding program could also facilitate the improvement in the nutritional quality of milk.

## Acknowledgments

This research received a financial support from the European Commission, Directorate-General for Agriculture and Rural Development, under Grant Agreement 211708 and from the Commission of the European Communities, FP7, KBBE-2007-1. This paper does not necessarily reflect the view of these institutions and in no way anticipates the Commission's future policy in this area. Hélène Soyeurt, a Post-doctoral Researcher, Jérémie Vandenplas, a Research Fellow and Nicolas Gengler, a Senior Research Associate until December 31, 2011, acknowledge also the support of the National Fund for Scientific Research (Brussels, Belgium). The authors also acknowledge the support of the National Fund for Scientific Research for its grant FRFC 2.4604.11. The Ministry of Agriculture of the Walloon Region of Belgium [Service public de Wallonie, Direction générale de l'Agriculture, des Ressources naturelles et de l'Environnement, Direction de la Recherche] is acknowledged for their financial support through the project NovaUdderHealth (D31-1207). The authors acknowledge the technical support provided by the Walloon Breeding Association (Ciney, Belgium), the milk laboratory (Comité du Lait, Battice, Belgium) and Sylvie Herbage who realizes the LTF quantification of the Walloon samples used for the validation. Scottish Agricultural College receives funding from the Scottish

Government and they are acknowledged for funding the long-term selection experiment producing milk samples used in this analysis. Joel Amdall at the AgSource Laboratory (Menomonie, WI, USA) is also acknowledged for providing MIR spectral data for cows in the UW–Madison herd. OSAM platform and LAECEA project (2972, Direction générale opérationnelle Agriculture – Ressources naturelles environnement, Direction du développement et de la vulgarization, Namur, Belgium) are also acknowledged for providing the Walloon mastitis data.

## References

- Baker HM and Baker EN 2004. Lactoferrin and Iron: structural and dynamic aspects of binding and release. *BioMetals* 17, 209–2016.
- Baker EN 2005. Lactoferrin: a multi-tasking protein par excellence. *Cellular and Molecular Life Sciences* 62, 2529–2530.
- Barbano DM and Dellavalle ME 1987. Rapid method for determination of milk casein content by infrared analysis. *Journal of Dairy Science* 70, 1524–1528.
- Baumrucker CR 2000. Mammary mechanisms for lactoferrin: interactions with IGFBP-3. *Biotechnology Agronomy, Society and Environment* 4, 5–12.
- Baumrucker CR, Schanbacher F, Shang Y and Green MH 2005. Lactoferrin interaction with retinoid signaling: cell growth and apoptosis in mammary cells. *Domestic Animal Endocrinology* 30, 289–303.
- Baveye S, Elass E, Mazurier J, Spik G and Legrand D 1999. Lactoferrin: a multifunctional glycoprotein involved in the modulation of the inflammatory process. *Clinical Chemistry and Laboratory Medicine* 37, 281–296.
- Coffey MP, Simm G, Oldham JD, Hill WG and Brotherstone S 2004. Genotype and diet effects on energy balance in the first three lactations of dairy cows. *Journal of Dairy Science* 87, 4318–4326.
- Coleman J, Pierce KM, Berry DP, Brennan A and Horan B 2009. The influence of genetic selection and feed system on the reproductive performance of spring-calving dairy cows within future pasture-based production systems. *Journal of Dairy Science* 92, 5258–5269.
- Dal Zotto R, De Marchi M, Cecchinato A, Penasa M, Cdoro M, Carnier P, Gallo L and Bittante G 2008. Reproducibility and repeatability of measures of milk coagulation properties and predictive ability of mid-infrared reflectance spectroscopy. *Journal of Dairy Science* 91, 4103–4112.
- Emanuelson U, Danell B and Philipsson J 1988. Genetic parameters for clinical mastitis, somatic cell counts, and milk production estimated by multiple-trait restricted maximum likelihood. *Journal of Dairy Science* 71, 467–476.
- Farnaud S and Evans RW 2003. Lactoferrin – a multifunctional protein with antimicrobial properties. *Molecular Immunology* 40, 395–405.
- Gaunt SN, Raffio N, Kingsbury ET, Damon RA, Johnson WH and Mitchell BA 1980. Variation of lactoferrin and mastitis and their heritabilities. *Journal of Dairy Science* 63, 1874–1880.
- Hagiwara S-I, Kawai K, Anri A and Nagahata H 2003. Lactoferrin concentrations in milk from normal and subclinical mastitis cows. *Journal of Veterinary Medical Science* 65, 319–323.
- Harmon RJ 1994. Physiology of mastitis and factors affecting somatic cell counts. *Journal of Dairy Science* 77, 2103–2112.
- Hruschka WR 1987. Data analysis: wavelength selection methods. In *Near-infrared technology in the agricultural and food industries* (ed. P Williams and K Norris), pp. 35–55. American Association of Cereal Chemists, St Paul, MN, USA.
- Kutilla T, Suojala L, Lehtolainen T, Saloniemä H, Kaartinen L, Tähti M, Seppälä K and Pyörälä S 2004. The efficacy of bovine lactoferrin in the treatment of cows with experimentally induced *Escherichia coli* mastitis. *Journal of Veterinary Pharmacology and Therapeutics* 27, 197–202.
- Lindmark-Mansson H, Bränning C, Aldén G and Paulsson M 2006. Relationship between somatic cell count, individual leukocyte populations and milk components in bovine udder quarter milk. *International Dairy Journal* 16, 717–727.
- Mead PE and Tweedie JW 1990. cDNA and protein sequence of bovine lactoferrin. *Nucleic Acids Research* 18, 7167.
- Pösö J and Mäntysaari EA 1996. Relationships between clinical mastitis, somatic cell score, and production for the first three lactations of Finnish Ayrshire. *Journal of Dairy Science* 79, 1284–1291.
- Prendiville R, Lewis E, Pierce KM and Buckley F 2010. Comparative grazing behavior of lactating Holstein–Friesian, Jersey, and Jersey × Holstein–Friesian dairy cows and its association with intake capacity and production efficiency. *Journal of Dairy Science* 93, 764–774.
- Pugovel G, Baumrucker CR, Sauerwein H, Rühl R, Ontsouka E, Hammon HM and Blum JM 2005. Effects of an enhanced vitamin A intake during the dry period on retinoids, lactoferrin, IGF system, mammary gland epithelial cell apoptosis, and subsequent lactation in dairy cows. *Journal of Dairy Science* 88, 1785–1800.
- Rutten MJM, Bovenhuis H, Hettinga KA, Van Vanlenberg HJF and Van Arendonck JAM 2009. Predicting bovine milk fat composition using infrared spectroscopy based on milk samples collected in winter and summer. *Journal of Dairy Science* 92, 6202–6209.
- SAS 1999. SAS/STAT user's guide. Version 8. SAS Institute Inc., Cary, NC, USA. 3884p.
- Sinnaeve G, Dardenne P, Agneessens R and Biston R 1994. The use of near infrared spectroscopy for the analysis of fresh grass silage. *Journal of Near Infrared Spectroscopy* 2, 79–84.
- Sivakesava S and Irudayaraj J 2002. Rapid determination of tetracycline in milk by FT-MIR and FT-NIR Spectroscopy. *Journal of Dairy Science* 85, 487–493.
- Sorensen LK, Lund M and Juul B 2003. Accuracy of Fourier transform infrared spectrometry in determination of casein in dairy cows' milk. *Journal of Dairy Research* 70, 445–452.
- Soyeurt H, Dardenne P, Lognay G, Veselko D, Mayeres P and Gengler N 2006. Estimating fatty acid content in cow milk using mid-infrared spectrometry. *Journal of Dairy Science* 89, 3690–3695.
- Soyeurt H, Colinet FG, Arnould VM-R, Dardenne P, Bertozzi C, Renaville R, Portetelle D and Gengler N 2007. Genetic variability of lactoferrin content estimated by mid-infrared spectrometry in bovine milk. *Journal of Dairy Science* 90, 4443–4450.
- Soyeurt H, Dehareng F, Gengler N, McParland S, Wall E, Berry DP and Coffey M 2011. Mid-infrared prediction of bovine milk fatty acids across multiple breeds, production systems, and countries. *Journal of Dairy Science* 94, 1657–1667.
- Tsuda H, Sekine K, Ushida Y, Kuhara T, Takasuka N, Iigo M, Seok Han B and Moore MA 2000. Milk and dairy products in cancer prevention: focus on bovine lactoferrin. *Mutation Research* 462, 227–233.
- Ward PP, Paz A and Conneely OM 2005. Multifunctional roles of lactoferrin: a critical overview. *Cellular and Molecular Life Science* 62, 2540–2548.
- Westerhaus MO 1990. Improving repeatability of NIR calibrations across instruments. In *Proceedings of the 3rd International Conference on Near Infrared Spectroscopy*, Brussels, Belgium, pp. 671–674. Agricultural Research Centre, Gembloux, Belgium.
- William P and Norris K 2001. Near-infrared technology in the agricultural and food industries. American Association of Cereal Chemists, St Paul, MN.
- Williams P 2007. Near-infrared technology – getting the best out of light. PDK Grain, Nanaimo, Canada.