



## Short communication

# Linseed oil presents different patterns of oxidation in real-time and accelerated aging assays



Caroline Douny<sup>a,\*</sup>, Rina Razanakolona<sup>b</sup>, Laurence Ribonnet<sup>b</sup>, Jérôme Milet<sup>c</sup>, Vincent Baeten<sup>d</sup>, Hervé Rogez<sup>e</sup>, Marie-Louise Scippo<sup>a</sup>, Yvan Larondelle<sup>b</sup>

<sup>a</sup> Department of Food Sciences, Laboratory of Food Analysis, FARAH – Veterinary Public Health, University of Liège, B43bis, Quartier Vallée 2, Avenue de Cureghem 10, 4000 Liège, Belgium

<sup>b</sup> Institut des Sciences de la Vie, Université catholique de Louvain, Croix du Sud, 2/L7.05.08, B-1348 Louvain-la-Neuve, Belgium

<sup>c</sup> Oxylent SA, Chemin du Fundus, 15, B-7822 Ghislenghien, Belgium

<sup>d</sup> Valorisation of Agricultural Products Department, Henseval Building, Walloon Agricultural Research Centre (CRA-W), 24 Chaussée de Namur, B-5030 Gembloux, Belgium

<sup>e</sup> Centre for Agro-Food Valorisation of Amazonian Bioactive Compounds (CVACBA), Universidade Federal do Para, 66.095-780 Belém, PA, Brazil

## ARTICLE INFO

## Article history:

Received 20 January 2016

Received in revised form 25 March 2016

Accepted 4 April 2016

Available online 4 April 2016

## Keywords:

Linseed oil

Accelerated aging

Schaal oven stability test

Real-time aging

Oxidation products

Myricetin

Antioxidant

## ABSTRACT

This study aimed at verifying if the hypothesis that one day at 60 °C is equivalent to one month at 20 °C could be confirmed during linseed oil aging for 6 months at 20 °C and 6 days at 60 °C using the “Schaal oven stability test”. Tests were conducted with linseed oil supplemented or not with myricetin or butyl-hydroxytoluene as antioxidants. Oxidation was evaluated with the peroxide and *p*-anisidine values, as well as the content in conjugated dienes and aldehydes. All four indicators of oxidation showed very different kinetic behaviors at 20 and 60 °C. The hypothesis is thus not verified for linseed oil, supplemented or not with antioxidant. In the control oil, the conjugated dienes and the peroxide value observed were respectively of 41.8 ± 0.8 Absorbance Unit (AU)/g oil and 254.3 ± 5.8 meq.O<sub>2</sub>/kg oil after 6 months at 20 °C. These values were of 18.2 ± 1.3 AU/g oil and 65.2 ± 20.3 meq.O<sub>2</sub>/kg after 6 days at 60 °C.

© 2016 Elsevier Ltd. All rights reserved.

## 1. Introduction

Linseed (*Linum usitatissimum*) oil has a high  $\alpha$ -linolenic acid (ALA) content and is therefore interesting in nutrition. Indeed, the consumption of n-3 polyunsaturated fatty acids (PUFA) is necessary for various physiological reasons and has been associated with a decrease of the incidence of inflammatory and cardiovascular pathologies (Simopoulos, 2008). Unfortunately, the high PUFA content of linseed oil also contributes to its rapid oxidation (Guillén & Uriarte, 2012). To prevent it, supplementation with antioxidants is required. Synthetic antioxidants commonly used in food include butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethoxyquin and ascorbyl palmitate (Reische, Lillard, & Eitenmiller, 2008). These antioxidants show an effective protection of the PUFA (Anwar, Siddiq, Iqbal, & Asi, 2007), but seri-

ous doubts concerning their safety oriented research towards the exploration of natural antioxidants (Chirinos, Huaman, Betalleluz-Pallardel, Pedreshi, & Campos, 2011; Martín et al., 2014), such as flavonoids, tocopherols and carotenoids (Pelli & Lyly, 2003). Michotte et al. (2011) showed that myricetin, a flavonol present in many plants, is an effective protector against autoxidation of PUFA in linseed oil. Other studies demonstrated that myricetin can also protect sunflower (Marinova, Toneva, & Yanishlieva, 2008) and rapeseed oils (Chen, Chan, Ho, Fung, & Wang, 1996).

The AOCS Cg 5-97 method, also known as the *Schaal oven stability test*, is an oven storage test used to simulate the real-time aging of oils (Michotte et al., 2011). This test should be conducted with oils as little oxidized as possible, at 60 °C in the dark. This method shows a good correlation with real-time stability studies. In the literature, some authors showed that the oxidation parameters obtained from aging realized at 60 °C and at a temperature close to ambient temperature are linearly linked. Indeed, Abou-Gharbia, Shehata, Youssef, and Shahidi (1996) indicated that when Tehina, a paste obtained with ground sesame seeds, was aged 2 days at 65 °C, the *p*-anisidine (PA) value and peroxide value

\* Corresponding author.

E-mail addresses: [cdouny@ulg.ac.be](mailto:cdouny@ulg.ac.be) (C. Douny), [rina.razanakolona@gmail.com](mailto:rina.razanakolona@gmail.com) (R. Razanakolona), [laurenceribonnet@hotmail.com](mailto:laurenceribonnet@hotmail.com) (L. Ribonnet), [Jerome.milet@oxylent.be](mailto:Jerome.milet@oxylent.be) (J. Milet), [V.baeten@cra.wallonie.be](mailto:V.baeten@cra.wallonie.be) (V. Baeten), [herverogez@gmail.com](mailto:herverogez@gmail.com) (H. Rogez), [miscippo@ulg.ac.be](mailto:miscippo@ulg.ac.be) (M.-L. Scippo), [yvan.larondelle@uclouvain.be](mailto:yvan.larondelle@uclouvain.be) (Y. Larondelle).

(PV) were similar to the values obtained with Tehina aged for 60 days at 22 °C. Similarly, soybean and cotton oils aged in closed bottles had equivalent «flavor scores» after 4 days at 60 °C and 4 months at 26 °C (Evans, List, Moser, & Cowan, 1973). Those works were reported by Khan and Shahidi (2001) and led to the hypothesis that an aging of 1 day at 65 °C was equivalent to an aging of 1 month at 25 °C. A linear relationship would mean that the results obtained at 60 °C during a short time experiment could be extrapolated to real-time (room temperature) aging of oil. Therefore, the objective of our study was to verify this hypothesis during linseed oil aging, without added antioxidant and with BHT or myricetin. In each aging experiment, primary products of PUFA oxidation were evaluated with the determination of conjugated dienes and PV; secondary products were evaluated with the determination of the PA value and aldehyde concentrations.

## 2. Materials and methods

### 2.1. Standards and reagents

1,1,3,3-tetraethoxypropane, 2,4-dinitrophenyl-hydrazine (97%), 2,4-decadienal, BHT (99%), PA were purchased from Sigma-Aldrich (St. Louis, Missouri). Stock solutions of 4-hydroxy-2-hexenal (4-HHE) and 4-hydroxy-2-nonenal (4-HNE) were obtained from Cayman Chemicals (Ann Arbor, Michigan). Myricetin was obtained from Extrasynthese (Genay, France).

Chloroform and water were of Chromanorm quality, while acetic acid 100% was of Normapur Quality, all provided by VWR International (Radnor, Pennsylvania). Ethanol absolute, trichloroacetic acid and hydrochloric acid 12 M were from Merck (Darmstadt, Germany). LC-MS-grade acetonitrile was obtained from Biosolve (Valkenswaard, The Netherlands).

### 2.2. Sample preparation and oxidation conditions

Refined linseed oil (RLO) free of synthetic antioxidant was kindly supplied by Vandeputte (Mouscron, Belgium). RLO was collected in a 1 L plastic bottle, inerted and stored at 4 °C until use. Samples of RLO (20 g), with added antioxidant and without (controls), were prepared as described by Michotte et al. (2011): BHT and myricetin were solubilized in a limited quantity of ethanol so that it did not exceed 4% of the weight of the final solution. Then, RLO was added in order to obtain the desired concentration of 555 µmol antioxidant/kg oil for each compound. The solution was strongly mixed with a glass rod for 10 min and flushed with nitrogen for 3 min to remove ethanol. The different mixtures were stored at 4 °C until the start of the experiments. The mixtures were then strongly shaken and distributed (20 g) in left open tinted glass bottles (100 ml, diam. 4 cm), which were stored for 6 days at 60 °C (accelerated aging) or 6 months at 20 °C (real-time aging) in two ventilated ovens (ULM800, Memmert, Schwabach, Germany). Both oven temperatures were monitored during the test with temperature probe from VWR International (Radnor, Pennsylvania). For the 60 °C condition, bottles were removed from the oven every 12 h for the two first days, then every 24 h until 6 days. For the 20 °C condition, bottles were removed from the oven every 2 weeks for the two first months, then every month until 6 months. Then, bottles were flushed with nitrogen, closed and stored at –70 °C until random analysis.

### 2.3. Conjugated dienes

Conjugated dienes were evaluated in 25 mg RLO diluted with 100 mL hexane by measuring the absorbance at 232 nm with a

UV-vis Cecil 2041 spectrophotometer (Cecil Instruments Limited, Cambridge, UK) (Dieffenbacher, 1992).

### 2.4. Peroxide value

The hydroperoxides were measured using the iodometric titration method (AOAC, 1995). Five g of RLO were diluted in 30 mL acetic acid/chloroform solution (3/2, v/v) then 0.5 mL of a saturated potassium iodide solution were added and a titration was realized with a 0.1 N Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution.

### 2.5. *p*-anisidine value

The PA value was determined by diluting 0.5 g RLO with 25 mL isooctane and adding PA (0.25% in glacial acetic acid) (American Oil Chemists' Society, 1998). PA value was measured at 350 nm.

### 2.6. Aldehyde measurement by LC-MS/MS

The aldehyde measurement was performed according to Douny et al. (2015). Briefly, 2 g of oil, added with BHT and methylmalondialdehyde as internal standard were extracted two times with water/ethanol 50/50 (v/v). Dinitrophenylhydrazone derivatives were prepared by adding a 2,4-dinitrophenyl-hydrazine solution (0.05 M in ethanol/HCl 12 M 9:1 (v/v)) and incubating for 2 h at 60 °C.

Separation and detection of aldehydes as derivatives were performed using a Spectra System P4000 HPLC system and a LCQ Deca ion trap mass spectrometer, with an Electrospray source (ThermoQuest Finnigan, San Jose, California).

### 2.7. Statistical analysis

Statistical Analysis System (SAS Institute, 2000) was used to check the data for normal distribution with a Shapiro–Wilk test and when variables were not normally distributed, a logarithmic transformation was performed. Significant differences between day 0 and other days/months ( $p < 0.05$ ) were tested using the general linear model procedure. The non-parametric test of Kruskal Wallis has been used when normality was not respected despite a logarithmic transformation.


## 3. Results

The determination of oxidation products obtained from RLO with and without added antioxidants aged at a temperature of 60 °C or 20 °C are presented in Fig. 1. Conjugated dienes (Fig. 1I and II), PV (Fig. 1III and IV) and PA value (Fig. 1V and VI) are expressed as mean ± standard deviation (SD) of three independent experiments, with measurements performed twice on each sample, leading to a total of 6 values per condition. MDA content (Fig. 1VII and VIII) is expressed as the mean concentration ± SD of three independent experiments, with measurements performed once on each sample, leading to a total of 3 values per condition.

### 3.1. Primary oxidation products

#### 3.1.1. Conjugated dienes

For the dienes, the absorbance value reached  $41.8 \pm 0.8$  AU/g oil after 6 months at 20 °C, for the control, and  $28.0 \pm 0.6$  AU/g oil, for the RLO samples supplemented with BHT, corresponding to an increase of respectively nearly 150% and 75%, compared to the absorbance value measured at day 0 ( $16.4 \pm 0.2$  AU/g oil for the control RLO and  $15.9 \pm 0.6$  AU/g oil for RLO supplemented with BHT) (Fig. 1.I). At this temperature, conjugated diene values



**Fig. 1.** Evolution of the products of oxidation expressed as conjugated dienes (CD, AU<sub>232nm</sub>/g oil) (I, II), peroxide value (PV, mequiv.O<sub>2</sub>/kg oil) (III, IV), p-anisidine value (p-AV, AU<sub>550nm</sub>/g oil) (V, VI) and malondialdehyde (MDA) (VII, VIII) concentration (μmol/kg oil) in refined linseed oil during real-time aging at 20 °C for 6 months (I, III, V, VII) or accelerated aging at 60 °C for 6 days (II,IV,VI,VIII). Mean ± standard deviation (SD) of three independent experiments. Significant differences comparing day 0 to other times of storage are indicated by a letter ( $P < 0.05$ ): a for control, b for BHT and c for myricetin. AU stands for Absorbance Unit.

became significantly different from those at day 0 ( $p < 0.05$ ) after 1.5 month of storage for both control and RLO containing BHT, while no significant difference compared to day 0 was recorded

for RLO containing myricetin (Fig. 1.I). At 60 °C, the absorbance value reached  $18.2 \pm 1.3$  AU/g oil for the control aged 6 days and  $17.5 \pm 0.5$  AU/g oil for the samples containing BHT, corresponding

to an increase of respectively nearly 50% and 43% compared to the absorbance value measured at day 0 ( $12.2 \pm 0.4$  AU/g oil for the control RLO and  $12.2 \pm 0.2$  AU/g oil for RLO supplemented with BHT) (Fig. 1II). At 60 °C, conjugated diene values became significantly different from those at day 0 ( $p < 0.05$ ) after 4 days of storage for both control RLO and RLO containing BHT, while, as it was observed at 20 °C, no significant difference compared to day 0 was recorded for RLO containing myricetin (Fig. 1II).

### 3.1.2. Peroxide value

At 20 °C, PV increased drastically in control RLO from  $1.8 \pm 0.3$  meq.O<sub>2</sub>/kg oil at day 0 to  $254.3 \pm 5.8$  meq.O<sub>2</sub>/kg oil after 6 months (Fig. 1III). In RLO samples supplemented with BHT or myricetin, the PV, after 6 months at 20 °C, was respectively  $113.2 \pm 3.4$  and  $18.9 \pm 1.7$  meq.O<sub>2</sub>/kg oil. At this temperature, PV became significantly different from those at day 0 ( $p < 0.05$ ) after 1 month of storage for both control and RLO containing BHT, and 1.5 month in RLO containing myricetin (Fig. 1III). After 6 days of storage at 60 °C, PV increased from  $2.4 \pm 1.5$  meq.O<sub>2</sub>/kg oil at day 0 to  $65.2 \pm 20.3$  meq.O<sub>2</sub>/kg oil in RLO without any added antioxidant, and to  $57.2 \pm 8.9$  and  $26.4 \pm 3.6$  meq.O<sub>2</sub>/kg oil in RLO samples containing BHT and myricetin, respectively (Fig. 1IV). After 1.5 day of storage at 60 °C, PV became significantly different from those at day 0 ( $p < 0.05$ ) for the control RLO, as well as for RLO containing BHT or myricetin (Fig. 1IV).

## 3.2. Secondary oxidation products

### 3.2.1. *p*-anisidine value

At 20 °C, after 6 months of storage, PA values reached  $23.7 \pm 8.1$  AU/g oil and  $10.3 \pm 0.8$  AU/g oil in control and BHT supplemented RLO, respectively, while it remained close to the value measured at day 0 (i.e.  $2.2 \pm 0.5$  AU/g oil) in RLO supplemented with myricetin (Fig. 1V). Compared to day 0, the PA values were significantly higher ( $p < 0.05$ ) in both the control and the RLO containing BHT from three and six months of storage at 20 °C, respectively (Fig. 1V). At 60 °C, a significant accumulation of aldehydes could be observed after 1.5 day of storage when looking at the PA values for the control and the RLO supplemented with BHT, and after 4 days of storage for the RLO containing myricetin (Fig. 1VI). The observed values increased from  $1.1 \pm 0.5$  AU/g oil (at day 0) to  $38.3 \pm 17.3$  AU/g oil for the control and to  $34.7 \pm 9.9$  and  $10.2 \pm 1.9$  AU/g oil for the RLO containing BHT and myricetin, respectively, after 6 days of accelerated aging.

### 3.2.2. Aldehyde levels

At 20 °C, in the control, the MDA content increased from  $5.6 \pm 1.8$  μmol/kg oil (day 0) to  $21.1 \pm 4.6$  μmol/kg oil after 6 months of storage (Fig. 1VII). MDA level became significantly different ( $p < 0.05$ ) from day 0 after 4 months of storage for the control while no significant difference compared to day 0 was observed for the RLO containing BHT or myricetin (Fig. 1VII).

At 60 °C, for the control, the MDA concentration increased from  $7.3 \pm 2.6$  μmol/kg oil (day 0) to  $61.5 \pm 6.8$  μmol/kg oil after 6 days of storage (Fig. 1VIII). As it was observed at 20 °C, a significant increase of MDA level was only observed in the control after 0.5 day of storage, while no significant difference compared to day 0 was observed in the RLO supplemented with BHT or myricetin (Fig. 1VIII).

Concerning the other aldehydes analyzed, 4-HNE and 2,4-decadienal were not detected in any sample at any time of aging, while 4-HHE was sporadically detected but always remained below the limit of quantification of the method (i.e.  $6.1$  μmol/kg oil).

## 4. Discussion

Without antioxidant, the conjugated dienes (Fig. 1I and II) and PV (Fig. 1III and IV) in RLO presented different patterns of increase, depending on the temperature. Indeed, they increased in a linear way at 60 °C while their increase was exponential at 20 °C. Moreover, the values obtained for these two parameters were nearly 3 times higher after 6 months at 20 °C than after 6 days at 60 °C, while the opposite trend has been observed for secondary oxidation products with aldehyde contents higher in RLO stored at 60 °C than in RLO stored at 20 °C. These observations suggest that the primary oxidation products accumulate in oil during real-time aging while they are transformed into other compounds during accelerated aging, mainly secondary oxidation products. As mentioned by Vieira and Regitano D'Arce (2001), conjugated dienes and hydroperoxides are very unstable at high temperature. They are then able to interact with other compounds (Cho, Endo, Fujimoto, & Kaneda, 1989) and they are rapidly transformed into secondary oxidation products like MDA (Zacheo, Cappello, & Perrone, 1998). Regarding MDA levels in the control during the accelerated aging, MDA formation was clearly faster at 60 °C and the final concentration (after 6 months at 20 °C or 6 days at 60 °C) was higher in RLO stored at 60 °C.

Whatever the measured oxidation products, myricetin was always better to prevent RLO from oxidation than BHT, at both temperatures. Regarding BHT, after 6 days of storage at 60 °C, the values of parameters measured in RLO containing this antioxidant were similar to those measured in the control. In contrast, during the aging experiment at 20 °C, the values of parameters measured in RLO containing BHT were lower than those observed for the control but higher than for the oil with myricetin. These findings could be explained by the fact that BHT could lose its antioxidant activity by reacting with secondary oxidation products (Chirinos et al., 2011). Concerning myricetin, this study confirms its high protective effect against lipid oxidation already shown in studies concerning ALA (Michotte et al., 2011), methyl linoleate (Pekkarinen, Heinonen, & Hopia, 1999) or lipid peroxidation in rat hepatocytes (Nuutila, Puupponen-Pimiä, Aarni, & Oksman-Caldentey, 2003).

## 5. Conclusion

When measuring PUFA oxidation, the difference in the behavior of RLO submitted to two different aging treatments, 60 °C or 20 °C, was visible after 3, 4 or 6 days or months of aging, respectively, depending on the oxidation products considered. It clearly appeared that the accelerated aging at 60 °C underestimates the primary oxidation products, as compared to a real-time aging at 20 °C, while the secondary products are overestimated at 60 °C.

In conclusion, the hypothesis that 1 day at 60 °C is equivalent to 1 month at 20 °C is not verified for RLO, with or without antioxidant.

## Conflict of interest

All others should have no conflict of interest.

This article does not contain any studies with human or animal subjects.

## Acknowledgements

Authors thanks the "Service public de Wallonie – DGO 6" for financial support and Nassim Moula (Department of Animal Production, FARAH, Université de Liège) for his skilled assistance in statistical analysis.



## References

- Abou-Gharbia, H. A., Shehata, A. A. Y., Youssef, M., & Shahidi, F. (1996). Oxidative stability of sesame paste (Tehina). *Journal of Food Lipids*, 3, 129–137.
- American Oil Chemists' Society (1998). *Official Methods and Recommended Practices of the American Oil Chemists' Society* (5th ed.). AOCS Press.
- Anwar, F., Siddiq, A., Iqbal, S., & Asi, M. R. (2007). Stabilization of sunflower oil with *Moringa oleifera* leaves under ambient storage. *Journal of Food Lipids*, 14, 35–49.
- AOAC Official Method 965.33 (1995). *Peroxide value of oils and fats. Titration method. Official methods of analysis, Oils and Fat*. Chapter 41, 9.
- Chen, Z. Y., Chan, P. T., Ho, K. Y., Fung, K. P., & Wang, J. (1996). Antioxydant activity of natural flavonoid is governed by the number and location of their aromatic hydroxyl groups. *Chemistry and Physics of Lipids*, 79, 157–163.
- Chirinos, R., Huaman, M., Betalleluz-Pallardel, I., Pedreshi, R., & Campos, D. (2011). Characterisation of phenolic compounds of Inca muna (*Clinopodium bolivianum*) leaves and the feasibility of their application to improve the oxidative stability of soybean during frying. *Food Chemistry*, 128, 711–716.
- Cho, S., Endo, K., Fujimoto, K., & Kaneda, T. (1989). Autoxidation of ethyl eicosapentaenoate in a defatted fish dry model system. *Bulletin of the Japanese Society of Scientific Fisheries*, 55, 545–552.
- Dieffenbacher, A. (1992). Standard methods for the analysis of oils, fats and derivatives. In *1st supplement to the 7th edition international union of pure and applied chemistry commission on oils, fats and derivatives*.
- Douny, C., Tihon, A., Bayonnet, P., Brose, F., Degand, G., Rozet, E., Milet, J., et al. (2015). Validation of the analytical procedure for the determination of malondialdehyde and three other aldehydes in vegetable oil using Liquid Chromatography coupled to tandem mass spectrometry (LC-MS/MS) and application to linseed oil. *Food Analytical Methods*, 8, 1425–1435.
- Evans, C. D., List, G. R., Moser, H. A., & Cowan, J. C. (1973). Long term storage of soybean and cottonseed salad oils. *Journal of the American Oil Chemists's Society*, 50, 218–222.
- Guillén, M. D., & Uriarte, P. S. (2012). Aldehydes contained in edible oils of a very different nature after prolonged heating at frying temperature: Presence of toxic oxygenated  $\alpha$ ,  $\beta$  unsaturated Aldehydes. *Food Chemistry*, 131, 915–926.
- Khan, M. A., & Shahidi, F. (2001). Effects of natural and synthetic antioxidants on the oxidative stability of borage and evening primrose triacylglycerols. *Food Chemistry*, 75, 431–437.
- Marinova, E., Toneva, A., & Yanishlieva, N. (2008). Synergistic antioxidant effect of  $\alpha$ -tocophérol and myricetin on the autoxidation of triacylglycerols of sunflower oil. *Food Chemistry*, 106, 628–633.
- Martín, J. M., Freire, P. F., Daimiel, L., Martínez-Botas, J., Sánchez, C. M., Lasunción, M. Á., Peropadre, A., et al. (2014). The antioxidant butylated hydroxyanisole potentiates the toxic effects of propylparaben in cultured mammalian cells. *Food and Chemical Toxicology*, 72, 195–203.
- Michotte, D., Rogez, H., Chirinos, R., Mignolet, E., Campos, D., & Larondelle, Y. (2011). Linseed oil stabilisation with pure natural phenolic compounds. *Food Chemistry*, 129, 1228–1231.
- Nuutila, A. M., Puupponen-Pimiä, R., Aarni, M., & Oksman-Caldentey, K. M. (2003). Comparison of antioxidant activities of onion and garlic extracts by inhibition of lipid peroxidation and radical scavenging activity. *Food Chemistry*, 81, 485–493.
- Pekkarinen, S. S., Heinonen, I. M., & Hopia, A. I. (1999). Flavonoids quercetin, myricetin, kaempferol and (+)-catechin as antioxidants in methyl linoleate. *Journal of the Science of Food and Agriculture*, 79, 499–506.
- Pelli, K., & Lyly, M. (2003). Les antioxydants dans l'alimentation. *INRA*, 28.
- Reische, D. W., Lillard, D. A., & Eitenmiller, R. R. (2008). Antioxydants. In C. C. Akoh & D. D. Min (Eds.), *Food lipids* (3rd ed., pp. 409–433). Taylor & Francis.
- Simopoulos, A. P. (2008). The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Experimental Biology and Medicine*, 233, 674–685.
- Vieira, T. M. F. S., & Regitano D'Arce, M. A. B. (2001). Canola oil thermal oxidation during oven test and microwave heating. *Lebensmittel-Wissenschaft und Technologie*, 34, 215–221.
- Zacheo, G., Cappello, L., & Perrone, L. M. (1998). Analysis of factors influencing lipid oxidation of almost seeds during accelerated ageing. *Lebensmittel-Wissenschaft und Technologie*, 31, 6–9.